



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/445, C07D 401/12	A1	(11) International Publication Number: WO 98/08513 (43) International Publication Date: 5 March 1998 (05.03.98)
(21) International Application Number: PCT/US97/14768 (22) International Filing Date: 22 August 1997 (22.08.97) (30) Priority Data: 60/024,831 28 August 1996 (28.08.96) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CUFF, George, W. [US/US]; 11902 Summerset Way South, Carmel, IN 46033 (US). THAKKAR, Arvind, L. [US/US]; 9020 Yellowwood Court, Indianapolis, IN 46260 (US). (74) Agents: STRODE, Janelle, D. et al.; Eli Lilly & Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: AMORPHOUS BENZOTHIOPHENES, METHODS OF PREPARATION, AND METHODS OF USE (57) Abstract The instant invention provides an amorphous form of a compound of formula (I). Methods of preparing the material, as well as methods of using same, are also provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

AMORPHOUS BENZOTHIOPHENES, METHODS OF PREPARATION,
AND METHODS OF USE

Osteoporosis describes a group of diseases which arises
5 from diverse etiologies, but which are characterized by the
net loss of bone mass per unit volume. The consequence of
this loss of bone mass and resulting bone fracture is the
failure of the skeleton to provide adequate support for the
body. One of the most common types of osteoporosis is
10 associated with menopause. Most women lose from about 20%
to about 60% of the bone mass in the trabecular compartment
of the bone within 3 to 6 years after the cessation of
menses. This rapid loss is generally associated with an
increase of bone resorption and formation. However, the
15 resorptive cycle is more dominant and the result is a net
loss of bone mass. Osteoporosis is a common and serious
disease among postmenopausal women.

There are an estimated 25 million women in the United
States alone who are afflicted with this disease. The
20 results of osteoporosis are personally harmful, and also
account for a large economic loss due to its chronicity and
the need for extensive and long term support
(hospitalization and nursing home care) from the disease
sequelae. This is especially true in more elderly patients.
25 Additionally, although osteoporosis is generally not thought
of as a life threatening condition, a 20% to 30% mortality
rate is related
to hip fractures in elderly women. A large percentage of
this mortality rate can be directly associated with
30 postmenopausal osteoporosis.

The most vulnerable tissue in the bone to the effects
of postmenopausal osteoporosis is the trabecular bone. This
tissue is often referred to as spongy or cancellous bone and
is particularly concentrated near the ends of the bone (near
35 the joints) and in the vertebrae of the spine. The
trabecular tissue is characterized by small osteoid
structures which interconnect with each other, as well as

-2-

the more solid and dense cortical tissue which makes up the outer surface and central shaft of the bone. This interconnected network of trabeculae gives lateral support to the outer cortical structure and is critical to the biomechanical strength of the overall structure. In postmenopausal osteoporosis, it is primarily the net resorption and loss of the trabeculae which leads to the failure and fracture of bone. In light of the loss of the trabeculae in the postmenopausal woman, it is not surprising that the most common fractures are those associated with bones which are highly dependent on trabecular support, for example, the vertebrae, the neck of the weight-bearing bones such as the femur and the forearm. Indeed, hip fracture, collics fractures, and vertebral crush fractures are hallmarks of postmenopausal osteoporosis.

The most generally accepted method for the treatment of postmenopausal osteoporosis is estrogen replacement therapy. Although therapy is generally successful, patient compliance with the therapy is low, primarily because estrogen treatment frequently produces undesirable side effects. An additional method of treatment would be the administration of a bisphosphonate compound, such as, for example, Fosamax® (Merck & Co., Inc.).

Throughout premenopausal time, most women have less incidence of cardiovascular disease than men of the same age. Following menopause, however, the rate of cardiovascular disease in women slowly increases to match the rate seen in men. This loss of protection has been linked to the loss of estrogen and, in particular, to the loss of estrogen's ability to regulate the levels of serum lipids. The nature of estrogen's ability to regulate serum lipids is not well understood, but evidence to date indicates that estrogen can up regulate the low density lipid (LDL) receptors in the liver to remove excess cholesterol. Additionally, estrogen appears to have some effect on the biosynthesis of cholesterol, and other beneficial effects on cardiovascular health.

-3-


It has been reported in the literature that serum lipid levels in postmenopausal women having estrogen replacement therapy return to concentrations found in the premenopausal state. Thus, estrogen would appear to be a reasonable
5 treatment for this condition. However, the side effects of estrogen replacement therapy are not acceptable to many women, thus limiting the use of this therapy. An ideal therapy for this condition would be an agent which regulates serum lipid levels in a manner analogous to estrogen, but
10 which is devoid of the side effects and risks associated with estrogen therapy.

A number of structurally unrelated compounds are capable of interacting with the estrogen receptor and producing unique in vivo profiles. Compounds with in vivo
15 profiles typical of a "pure" antagonist (for example, ICI 164,384) or of a relatively "pure" agonist (for example, 17 β -estradiol) represent opposite ends of a spectrum in this classification. Between these two extremes lie the SERMs ("selective estrogen receptor modulator"), characterized by
20 clinical and/or preclinical selectivity as full or partial agonists in certain desired tissues (for example, bone), and antagonists or minimal agonists in reproductive tissues. Within this pharmacologic class, individual SERMs may be further differentiated based on profiles of activity in
25 reproductive tissues. Raloxifene, a second generation SERM, displays potentially useful selectivity in uterine tissue with apparent advantages over triphenylethylene-based estrogen receptor ligands. As such, raloxifene appears to be well-suited at least for the treatment of postmenopausal
30 complications, including osteoporosis and cardiovascular disease. It is anticipated that, as further advances are made in the pharmacology and molecular biology of estrogen receptor active agents, further subclassifications of SERMs may evolve in the future along with an increased
35 understanding of the therapeutic utility of these novel classes of estrogenic compounds.

The advancement of raloxifene, in particular, has been somewhat hampered by its physical characteristics, both as to bioavailability and manufacturing. For example, raloxifene is generally insoluble, which may affect
5 bioavailability. Clearly, any improvement in the physical characteristics of raloxifene and in closely related compounds would potentially offer a more beneficial therapy and enhanced manufacturing capabilities.

Thus, it would be a significant contribution to the art to
10 provide amorphous forms of raloxifene and related compounds which have increased solubility, methods of preparation, pharmaceutical formulations, and methods of use.

The present invention provides a compound of formula I
15



20 wherein:

R¹ and R³ are independently hydrogen, -CH₃, -CO(C₁-C₆ alkyl), or -COAr, wherein Ar is optionally substituted phenyl;

R² is selected from the group consisting of
25 pyrrolidinyl, hexamethyleneimino, and piperidinyl; or a pharmaceutically acceptable salt or solvate thereof, in an amorphous form.

Figure 1 illustrates the mean raloxifene levels in plasma from dogs treated with either crystalline or amorphous forms of a compound of formula I.

5 The present invention further provides pharmaceutical formulations containing a compound of formula I.

Still further provided by way of the instant invention are processes for the preparation of amorphous forms of compounds of formula I.

10 The instant invention also provides methods of use for the compounds of formula I, including the inhibition of bone loss or bone resorption.

General terms used in the description of compounds herein described bear their usual meanings. For example, 15 the term "C₁-C₆ alkyl" represents a straight or branched alkyl chain having from 1 to 6 carbon atoms. Typical C₁-C₆ alkyl groups include methyl, ethyl, *n*-propyl, and *n*-butyl. The term "C₁-C₄ alkoxy" represents groups such as methoxy, ethoxy, *n*-propoxy, and *n*-butoxy.

20 Optionally substituted phenyl includes phenyl and phenyl substituted with C₁-C₆ alkyl, C₁-C₄ alkoxy, hydroxy, nitro, chloro, fluoro, or tri(chloro or fluoro)methyl, and the like.

The term "inhibit" includes its generally accepted 25 meaning which includes prohibiting, preventing, restraining, and slowing, stopping, or reversing progression, severity, or ameliorating a resultant symptom or effect.

Preferred embodiments of the current invention are compounds of formula I which include compounds wherein R¹ 30 and R³ are hydrogen, and R² is pyrrolidinyl, piperidinyl, or a hexamethyleneimino group. Representative compounds from this preferred group include:

6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-
pyrrolidinylethoxy)benzoyl]benzo[b]-thiophene,
35 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinyl-
ethoxy)benzoyl]benzo[b]thiophene, and

6-hydroxy-2-(4-hydroxy-phenyl)-3-[4-(2-hexamethyleneiminoethoxy)benzoyl]benzo[b]-thiophene.

More preferably, the product of the present invention is the Formula I compounds wherein R² is a piperidinyl group. A representative compound from this more preferred group includes: 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene.

The compounds of the current invention may be made according to established procedures, such as those detailed in U.S. Patent Nos. 4,133,814, 4,418,068, and 4,380,635, and European Patent Application 95306050.6, Publication No. 0699672, Kjell, et al., filed August 30, 1995, published March 6, 1996, all of which are herein incorporated by reference. In addition, the information disclosed in the published European Patent Application number 0670162 A1, published on September 6, 1995, is herein incorporated by reference. A crystalline form of raloxifene hydrochloride may be prepared by the methods disclosed in the Example section, *infra*.

The term "amorphous" includes a physical state which may be verified by x-ray diffraction and other means including but not limited to observation with a polarized light microscope and differential scanning calorimetry.

In general, the process starts with a benzo[b]thiophene having a 6-hydroxyl group and a 2-(4-hydroxyphenyl) group. The starting compound is protected, acylated, and deprotected to form the compounds of formula I. Further examples of the preparation of such compounds are also provided in the references discussed above.

The compound of formula I according to the present invention is conveniently prepared by a process which constitutes a further feature of the present invention, and which comprises recovering a compound of formula I from a solution thereof under conditions whereby an amorphous product is obtained.

The amorphous material of the instant invention was prepared by dissolving a crystalline form of a compound of

formula I in a suitable solvent or solvent mixture, such as, for example, methanol and water, followed by recovery of the material by any suitable means. Techniques which may be employed to recover amorphous compounds of formula I from the solution include those wherein the solvent is removed from the solution, preferably rapidly, and the product deposited, and those wherein the product is precipitated from a solution. Methods involving the use of these procedures which have been found to be satisfactory include spray drying, roller drying, solvent precipitation, rotary evaporation, and freeze drying. Particularly preferred for the practice of the present invention is the method of spray drying.

Solvents which may be employed in the practice of the present invention will be chosen according to the technique and conditions to be employed, and include water, methanol, ethanol, and the like, including mixtures thereof, if desired.

The concentration of a compound of formula I in the solvent is advantageously as high as possible, commensurate with an amorphous form of a compound of formula I being obtained, with preferable concentrations being in the range of from about 5 mg/ml to about 40 mg/ml. The higher concentrations obtained will typically depend on the solvent system employed in the preparation, and/or the presence or absence of povidone (PVP) or hydroxypropyl-b-cyclodextrin (HPBCD). The solvents may, if desired, be heated as an aid to solubility and solvent removal.

In general, the compounds of formula I have sufficient heat stability to withstand spray drying and the like, and accordingly, spray drying is the preferred method of recovery. Spray drying systems may be operated in a known manner to obtain an amorphous product essentially free from crystalline material as well as free from particulate contaminants. Closed cycle spray drying systems in which the drying medium is recycled are particularly safe and economic for use in obtaining the product of the present invention.

The drying gas employed in the process may be air, but preferred for the use with flammable solvents are inert gases

such as, for example, nitrogen, argon and carbon dioxide. Preferred would be nitrogen. The gas inlet temperature to the spray dryer is chosen according to the solvent employed, but would be, for example, in the range of from about 75 °C to about 150 °C.

The presence of the amorphous form of a compound of formula I was determined by observing the material under a polarized light microscope, and determining if the material was birefringent. If no birefringence was observed, the material was considered to be amorphous.

A compound of formula I in accordance with the present invention is preferably essentially free from the crystalline form of the material. Long term studies have indicated the amorphous form of the instant invention is very stable.

However, once recovered as amorphous material, conversion to the crystalline form may be prevented by the addition of any number of stabilizer materials known in the art, such as, for example, povidone, hydroxypropyl methylcellulose (HPMC), hydroxypropylcellulose (HPC), polyethylene glycol (PEG), hydroxypropyl- β -cyclodextrin (HPB), cyclodextrin, and the like.

Solubility of the amorphous form was demonstrated to be approximately 250 times greater than the crystalline form. Advantages of increased solubility include but are not limited to ease in processing the amorphous material, which includes equipment cleaning issues; ease in formulation and delivery of the material, and the like.

A compound of formula I in an amorphous form may also be combined with a number of other materials prior to or after spray drying, or otherwise processed to provide amorphous material, which may in turn be further formulated for processing.

Compounds of formula I which are amorphous have been demonstrated to have several advantages, including but not limited to a high degree of bioavailability, as well as being in a form for effective methods of administration.

The term "solvate" represents an aggregate that comprises one or more molecules of the solute, such as a formula I compound, with one or more molecules of solvent. Although the free-base form of formula I compounds can be used in the methods of the present invention, it is preferred to prepare and use a pharmaceutically acceptable salt form. The term "pharmaceutically acceptable salt" refers to either acid or base addition salts which are known to be non-toxic and are commonly used in the pharmaceutical literature. The pharmaceutically acceptable salts generally have enhanced solubility characteristics compared to the compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions. The compounds used in the methods of this invention primarily form pharmaceutically acceptable acid addition salts with a wide variety of organic and inorganic acids, and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Such salts are also part of this invention. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, hypophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, caproate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycolate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, terephthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate,

pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzenesulfonate, p-bromophenylsulfonate, chlorobenzenesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like. A preferred salt is the hydrochloride salt.

The pharmaceutically acceptable acid addition salts are typically formed by reacting a compound of formula I with an equimolar or excess amount of acid. The reactants are generally combined in a mutual solvent such as diethyl ether or ethyl acetate. The salt normally precipitates out of solution within about one hour to 10 days and can be isolated by filtration, or the solvent can be stripped off by conventional means. The present invention further provides for pharmaceutically acceptable formulations for administering to a mammal, including humans, in need of treatment, which comprises an effective amount of a compound of formula I and a pharmaceutically acceptable diluent or carrier.

As used herein, the term "effective amount" means an amount of compound of the present invention which is capable of inhibiting, alleviating, ameliorating, treating, or preventing further symptoms in mammals, including humans, suffering from estrogen deprivation, for example, menopause or ovariectomy, or inappropriate estrogen stimulation such as uterine fibrosis or endometriosis, or suffering from aortal smooth muscle cell proliferation or restenosis. In the case of estrogen-dependent cancers, the term "effective amount" means the amount of compound of the present invention which is capable of alleviating, ameliorating, inhibiting cancer growth, treating, or preventing the cancer and/or its symptoms in mammals, including humans.

By "pharmaceutically acceptable formulation" it is meant that the carrier, diluent, excipients and salt must be

-11-

compatible with the active ingredient (a compound of formula I) of the formulation, and not be deleterious to the recipient thereof. Pharmaceutical formulations can be prepared by procedures known in the art. For example, the compounds of this invention can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, and the like. Examples of excipients, diluents, and carriers that are suitable for such formulations include the following: fillers and extenders such as starch, sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as povidone, sodium starch glycolate, sodium carboxymethylcellulose, agar agar, calcium carbonate, and sodium bicarbonate; agents for retarding dissolution such as paraffin; resorption accelerators such as quaternary ammonium compounds; surface active agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as kaolin and bentonite; and lubricants such as talc, calcium and magnesium stearate and solid polyethylene glycols. Final pharmaceutical forms may be: pills, tablets, powders, lozenges, sachets, cachets, or sterile packaged powders, and the like, depending on the type of excipient used.

Additionally, the compounds of this invention are well suited to formulation as sustained release dosage forms. The formulations can also be so constituted that they release the active ingredient only or preferably in a particular part of the intestinal tract, possibly over a period of time. Such formulations would involve coatings, envelopes, or protective matrices which may be made from polymeric substances or waxes.

The particular dosage of a compound of formula I required to treat, inhibit, or prevent the symptoms and/ or disease of a mammal, including humans, suffering from the above maladies according to this invention will depend upon the particular disease, symptoms, and severity. Dosage,

-12-

routes of administration, and frequency of dosing is best decided by the attending physician. Generally, accepted and effective doses will be from 15mg to 1000mg, and more typically from 15mg to 80mg. Such dosages will be administered to a patient in need of treatment from one to three times each day or as often as needed for efficacy, and for periods of at least two months, more typically for at least six months, or chronically.

As a further embodiment of the invention, the compounds of formula I may be administered along with an effective amount of an additional therapeutic agent, including but not limited to estrogen, progestin, benzothiophene compounds having including raloxifene, naphthyl compounds having antiestrogen activity, bisphosphonate compounds such as alendronate and tiludronate, parathyroid hormone (PTH), including truncated and/or recombinant forms of PTH such as, for example, PTH (1-34), calcitonin, bone morphogenic proteins (BMPs), or combinations thereof. The different forms of these additional therapeutic agents available as well as the various utilities associated with same and the applicable dosing regimens are well known to those of skill in the art.

Various forms of estrogen and progestin are commercially available. As used herein, the term "estrogen" includes compounds having estrogen activity and estrogen-based agents. Estrogen compounds useful in the practice of the present invention include, for example, estradiol estrone, estriol, equilin, equilenin, estradiol cypionate, estradiol valerate, ethynyl estradiol, polyestradiol phosphate, estropipate, diethylstilbestrol, dienestrol, chlorotrianisene, and mixtures thereof. Estrogen-based agents, include, for example, 17-a-ethynyl estradiol (0.01-0.03 mg/day), mestranol (0.05-0.15 mg/day), and conjugated estrogenic hormones such as Premarin® (Wyeth-Ayerst; 0.2-2.5 mg/day). As used herein, the term "progestin" includes compounds having progestational activity such as, for example, progesterone, norethynodrel, norgestrel, megestrol

-13-

acetate, norethindrone, progestin-based agents, and the like. Progestin-based agents include, for example, medroxyprogesterone such as Provera® (Upjohn; 2.5-10 mg/day), norethynodrel (1.0-10.0 mg/day), and norethindrone (0.5-2.0 mg/day). A preferred estrogen-based compound is Premarin®, and norethynodrel and norethindrone are preferred progestin-based agents. The method of administration of each estrogen- and progestin-based agent is consistent with that known in the art.

The formulations which follow are given for purposes of illustration and are not intended to be limiting in any way. The total active ingredients in such formulations comprises from 0.1% to 99.9% by weight of the formulation. The term "active ingredient" means a compound of formula I.

Formulation 1: Gelatin Capsules

Ingredient	Quantity (mg/capsule)
Active Ingredient	0.1-1000
Starch NF	0-500
Starch flowable powder	0-500
Silicone fluid 350 centistokes	0-15

The ingredients are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules.

Formulation 2: Tablets

Ingredient	Quantity (mg/tablet)
Active Ingredient	2.5-1000
Starch	10-50
Cellulose, microcrystalline	10-20
Polyvinylpyrrolidone (as 10% solution in water)	5
Sodium carboxymethylcellulose	5
Magnesium stearate	1
Talc	1-5

-14-

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granules thus produced are dried at 50-60 °C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethylcellulose, magnesium stearate, and talc, previously passed through a No. 60 mesh U.S. sieve, are added to the above granules and thoroughly mixed. The resultant material is compressed in a tablet forming machine to yield the tablets.

Formulation 3: Suppositories

Ingredient	Weight
Active ingredient	150 mg
Saturated fatty acid glycerides	3000mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the fatty acid glycerides which had previously heated to their melting point. The mixture is poured into a suppository mold and allowed to cool.

The following examples and preparations are provided to better elucidate the practice of the present invention and should not be interpreted in any way as to limit the scope of same. Those skilled in the art will recognize that various modifications may be made while not departing from the spirit and scope of the invention. All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

NMR data for the following Examples were generated on a GE 300 MHz NMR instrument, and anhydrous d-6 DMSO was used as the solvent unless otherwise indicated.

All experiments were run under positive pressure of dry nitrogen. All solvents and reagents were used as obtained.

-15-

The percentages are generally calculated on a weight (w/w) basis; except for HPLC solvents which are calculated on a volume (v/v) basis. Proton nuclear magnetic resonance (^1H NMR) spectra were obtained on a Bruker AC-300 FTNMR spectrometer at 300.135 MHz. Melting points were determined by differential scanning calorimetry (DSC) in a TA Instrument DCS 2920 using a closed cell and a heating rate of 2°C/minute. The reactions were generally monitored for completion using high performance liquid chromatography (HPLC). The X-ray powder diffraction spectra were obtained in a Siemens D5000 X-Ray Powder Diffraktometer, using copper radiation and a Si(Li) detector. Some reactions were monitored using a Zorbax Rx-C8 column, (25 cm x 4.6 mm ID, 5 m particle) eluting with a mixture of 60 mM phosphate (KH_2PO_4) and 10 mM octanesulfonate (pH 2.0)/ acetonitrile (60:40).

The acylation, dealkylation, or acylation/dealkylation reactions are also monitored for completion by HPLC. A sample of the reaction mixture was assayed using a Zorbax Rx-C8 column, (25 cm x 4.6 mm ID, 5 m particle), eluting with a gradient as shown below:

GRADIENT SOLVENT SYSTEM

	Time (min.)	A (%)	B (%)
25	0	60	40
	5	60	40
	10	45	55
	20	38	62
	25	45	55
30	32	45	55
	37	60	40
	42	60	40

A: 0.05 M HClO_4 (pH=2.0)

35 B: acetonitrile

The reaction mixture was analyzed by diluting a 0.1 to

-16-

0.2 mL sample to 50 mL with a 60:40 mixture of A/B. Similarly, the mother liquor of the recrystallizations was sampled in a similar manner.

The amount (percentages) of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride in the crystalline material (potency) was determined by the following method. A sample of the crystalline solid (5 mg) was weighed into a 100-mL volumetric flask, and dissolved in a 70/30 (v/v) mixture of 75 mM potassium phosphate buffer (pH 2.0) and acetonitrile. An aliquot of this solution (10 mL) was assayed by high performance liquid chromatography, using a Zorbax Rx-C8 column (25 cm x 4.6 mm ID, 5 µm particle) and UV detection (280 nm). The following gradient solvent system is used:

Gradient Solvent System (Potency)

	time (min)	A (%)	B (%)
	0	70	30
	12	70	30
20	14	25	75
	16	70	30
	25	70	30

A: 75 mM KH₂PO₄ buffer (pH 2.0)

B: acetonitrile

The percentage of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride in the sample is calculated using the peak area, slope (m), and intercept (b) of the calibration curve with the following equation:

$$\% \text{ potency} = \frac{\text{peak area} - b}{m} \times \frac{\text{sample volume (mL)}}{\text{sample weight (mg)}}$$

The amount (percentage) of solvent, such as methanol, ethanol, or 1,2-dichloroethane, present in the crystalline material is determined by gas chromatography. A sample of

-17-

the crystalline solid (50 mg) was weighed into a 10-mL volumetric flask, and dissolved in a solution of 2-butanol (0.025 mg/mL) in dimethylsulfoxide. A sample of this solution was analyzed on a gas chromatograph using a DB Wax column (30 m x 0.53 mm ID, 1 m particle), with a column flow of 10 mL/min and flame ionization detection. The column temperature was heated from 35°C to 230°C over a 12 minute period. The amount of solvent was determined by comparison to the internal standard (2-butanol), using the following formula:

$$\% \text{ solvent} = \frac{C}{D} \times \frac{E}{F} \times \frac{G}{H} \times I$$

wherein:

- 15 C = ratio of solvent in sample
- D = average ratio of standard for specific solvent
- E = average weight of standard
- F = weight of sample (mg)
- G = volume of sample (10 mL)
- 20 H = volume of standard (10,000 mL)
- I = purity of standard (%)

Preparation 1

6-Methoxy-2-(4-methoxyphenyl)benzo[b]thiophene

25 A solution of 3-methoxybenzenethiol (100 grams) and potassium hydroxide (39.1 grams) in water (300 mL) was added to denatured ethanol (750 mL), and the resulting mixture cooled to about 0°C. The cold mixture was treated with 4'-methoxyphenacyl bromide (164 grams) in several small portions. Upon complete addition, the mixture was cooled for an additional ten minutes, then allowed to warm to room temperature. After three hours, the mixture was concentrated *in vacuo*, and the residue treated with water (200 mL). The resulting mixture was treated with ethyl acetate, and the phases were separated. The organic phase

-18-

was washed with water (2x), sodium bicarbonate solution (2x), and sodium chloride solution (2x). The organic phase was then dried over magnesium sulfate, filtered, and evaporated to dryness *in vacuo* to give 202 grams of α -(3-methoxyphenylthio)-4-methoxyacetophenone. This crude product was crystallized from methanol and washed with hexane to give 158 grams. Melting point 53°C.

Polyphosphoric acid (930 grams) was heated to 85°C and treated with the intermediate product from above (124 grams) in small portions over 30 minutes. Upon complete addition, the resulting mixture was stirred at 90°C. After an additional 45 minutes, the reaction mixture was allowed to cool to room temperature. This mixture was treated with crushed ice while the mixture was cooled in an ice bath. The resulting mixture was treated with water (100 mL) producing a light pink precipitate. The precipitate was isolated by filtration, washed with water and methanol, and dried *in vacuo* at 40°C to give 119 grams of 6-methoxy-2-(4-methoxyphenyl) benzo[b]thiophene. This crude product was slurried in hot methanol, filtered, and washed with cold methanol. The resulting solid material was recrystallized from ethyl acetate (4 liters), filtered, washed with hexane, and dried *in vacuo* to 68 grams of the title compound. Melting point 187-190.5°C.

Preparation 2

Ethyl 4-(2-Piperidinylethoxy)benzoate

A mixture of ethyl 4-hydroxybenzoate (8.31 g), 1-(2-chloroethyl)piperidine monohydrochloride (10.13 g), potassium carbonate (16.59 g), and methyl ethyl ketone (60 mL) was heated to 80°C. After one hour, the mixture was cooled to about 55°C and treated with additional 1-(2-chloroethyl)piperidine mono-hydrochloride (0.92 g). The resulting mixture was heated to 80°C. The reaction was monitored by thin layer chromatography (TLC), using silica-gel plates and ethyl acetate/acetonitrile/ triethylamine (10:6:1, v/v). Additional portions of 1-(2-

-19-

chloroethyl)piperidine hydrochloride are added until the starting 4-hydroxybenzoate ester is consumed. Upon complete reaction, the reaction mixture was treated with water (60 mL) and allowed to cool to room temperature. The aqueous layer was discarded and the organic layer concentrated *in vacuo* at 40°C and 40 mm Hg. The resulting oil was used in the next step without further purification.

Preparation 3

4-(2-Piperidinylethoxy)benzoic Acid Hydrochloride

10

A solution of the compound prepared as described in Preparation 2 (about 13.87 g) in methanol (30 mL) was treated with 5 N sodium hydroxide (15 mL), and heated to 40°C. After 4 1/2 hours, water (40 mL) was added. The resulting mixture was cooled to 5-10°C, and concentrated hydrochloric acid (18 mL) was added slowly. The title compound crystallized during acidification. This crystalline product was collected by filtration, and dried *in vacuo* at 40-50°C to give 83% yield of the title compound. Melting point 270-271°C.

Preparation 4

6-Methoxy-2-(4-methoxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]-benzo[b]thiophene Hydrochloride

25

A mixture of the compound prepared as described in Preparation 1 (8.46 grams) and the acid chloride prepared as described in Preparation 3 (10.0 grams) in methylene chloride (350 mL) was cooled to about 20-25°C. The cool mixture was treated with boron trichloride (2.6 mL), and the resulting mixture mechanically stirred. The reaction was monitored by HPLC using the assay described above. After 85 minutes, the *in situ* HPLC yield based on a 6-methoxy-2-(4-methoxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene standard was 88%.

30
35

-20-

Preparation 5

6-Hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]-benzo[b]thiophene Hydrochloride
1,2-Dichloroethane Solvate
(Crystal Form I)

A solution of 6-methoxy-2-(4-methoxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride (2.0 g) in 1,2-dichloroethane (20 mL) was treated with boron trichloride (2.0 mL). The resulting mixture was stirred at 35°C for about 18 hours. A mixture of ethanol and methanol (10 mL, 95:5, 3A) was treated with the reaction mixture from above, causing the alcoholic mixture to reflux. Upon complete addition, the resulting crystalline slurry was stirred at 25°C. After one hour, the crystalline product was filtered, washed with cold ethanol (10 mL), and dried at 40°C in vacuo to give 1.78 g of the title compound. The X-ray powder diffraction pattern is identical to that reported in Table 1.

Potency: 80.2%

1,2-Dichloroethane: 7.5% (gas chromatography)

Table 1. X-ray Diffraction Pattern for Crystal Form 1.

	d-line spacing (Angstroms)	I/I ₀ (x100)
25	16.1265	3.80
	10.3744	8.63
	8.3746	5.29
	7.9883	36.71
30	7.2701	5.06
	6.5567	70.77
	6.2531	6.79
	5.5616	24.05
	5.3879	100.00
35	5.0471	89.64
	4.7391	85.96
	4.6777	39.36

-21-

	4.6332	62.60
	4.5191	77.56
	4.2867	36.82
	4.2365	41.66
5	4.1816	49.60
	4.0900	11.28
	3.9496	11.85
	3.7869	36.25
	3.7577	56.16
10	3.6509	40.62
	3.5751	15.65
	3.5181	21.52
	3.4964	18.53
	3.4361	33.60
15	3.3610	6.21
	3.3115	4.95
	3.2564	7.36
	3.2002	3.80
	3.1199	15.77
20	3.0347	14.84
	2.8744	9.67
	2.8174	10.82
	2.7363	11.51

25 The amount of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-
piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride
present in the crystalline material is about 87.1%, as
determined using the high performance liquid chromatography
(HPLC) assay described below. The amount of 1,2-
30 dichloroethane present in the crystalline material is about
0.55 molar equivalents, as determined by proton nuclear
magnetic resonance spectroscopy.

-22-

Preparation 6

6-Hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinyloxy)benzoyl]-benzo[b]thiophene Hydrochloride
1,2-Dichloroethane Solvate
(Crystal Form I)

5

A mixture of the compound prepared as described in Preparation 2 (15 g) and dimethylformamide (0.2 mL) in 1,2-dichloroethane (250 mL) was cooled to 0°C. Phosgene (8.25 mL) was condensed in a cold, jacketed addition funnel (-10°C), and added to the cold mixture over a period of two minutes. The resulting mixture was heated to about 47°C. After about two and one half hours, the reaction was assayed by HPLC for completion. Additional phosgene may be added to drive the reaction to completion. Excess phosgene was removed by vacuum distillation at 30-32°C and 105-110 mm Hg.

10

After about three to four hours, the reaction solution was treated with the compound prepared as described in Preparation 1 (13.52 g). The resulting solution was cooled to 0°C. Boron trichloride (12.8 mL) was condensed in a graduated cylinder, and added to the cold reaction mixture. After eight hours at 0°C, the reaction solution was treated with additional boron trichloride (12.8 mL). The resulting solution was heated to 30°C. After 15 hours, the reaction was monitored for completion by HPLC.

15

A mixture of ethanol and methanol (125 mL, 95:5, 3A) was heated to reflux, and treated with the reaction solution from above over a 60 minute period. Upon complete addition, the acylation/demethylation reaction flask was rinsed with additional ethanol (30 mL). The resulting slurry was allowed to cool to room temperature with stirring. After one hour at room temperature, the crystalline product was filtered, washed with ethanol (75 mL), and dried at 40°C in vacuo to give 25.9 g of the title compound. The X-ray powder diffraction pattern is reported in Table 1. Melting point 261°C.

20

25

Potency: 87.1%

1,2-Dichloroethane: 0.55 molar equivalents (¹H NMR)

Preparation 7

6-Hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-
5 piperidinylethoxy)benzoyl]-benzo[b]thiophene Hydrochloride
Chlorobenzene Solvate
(Crystal Form 3)

A solution of the compound prepared as described in
Preparation 1 (2.92 grams) and the acid chloride prepared as
10 described in Preparation 4 (3.45 grams) in chlorobenzene (52
mL) was cooled to about 0°C. The cold solution was treated
with boron trichloride (2.8 mL). The resulting mixture was
mechanically stirred at about 0°C. After three hours,
15 additional boron trichloride (2.8 mL) was added, and the
reaction mixture was allowed to warm to room temperature.
After about 16-20 hours, the reaction mixture was cooled to
0°C. The cold reaction was quenched by the slow addition of
ethanol (26 mL) over 30 minutes. During the addition of the
20 alcohol, a crystalline solid formed. Upon complete addition
of the alcohol, the resulting mixture was stirred at room
temperature for one hour. The crystalline solid was removed
by filtration, washed with cold ethanol (25 mL), and dried
in vacuo at 40°C to give 5.94 grams of the title compound as
a yellow solid. The X-ray powder diffraction pattern is
25 identical to that reported in Table 2. Melting point
247°C.

Potency: 78.6%

Chlorobenzene: 12.3% (HPLC)

30 Table 2. X-ray Diffraction Pattern for Crystal Form III.

	d-line spacing	I/I ₀
	(Angstroms)	(x100)
	14.3518	7.24
	10.3335	6.17
35	8.8305	4.29
	7.9475	38.16

-24-

	6.5904	64.25
	6.2848	6.52
	5.6048	28.06
	5.4107	100.00
5	5.1544	11.26
	5.0493	53.26
	5.0224	46.11
	4.8330	76.94
	4.7694	34.23
10	4.6461	50.22
	4.5754	38.61
	4.4953	72.65
	4.3531	49.15
	4.2940	41.64
15	4.2425	35.75
	4.1856	21.63
	4.1338	9.47
	4.0793	12.69
	3.9960	18.50
20	3.9037	9.03
	3.7854	40.39
	3.7521	54.16
	3.6787	28.60
	3.6509	17.96
25	3.5444	31.72
	3.4679	41.55
	3.3899	7.69
	3.3101	5.72
	3.2561	7.42
30	3.1784	15.19
	3.0445	11.17
	3.0146	8.94
	2.9160	11.89
	2.8217	18.23
35	2.7500	12.06
	2.6436	9.65
	2.6156	6.97

The amount of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride present in the crystalline material is about 78.6%. The amount of chlorobenzene present in the crystalline material is about 12.3%, as determined by HPLC.

Preparation 8

6-Hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene Hydrochloride

A solution of sodium hydroxide (0.313 g) in methanol (10 mL) was diluted with additional methanol (40 mL) and water (10 mL). This solution was treated with the compound prepared as described in Example 5 (4.0 g). The resulting solution was extracted with hexane (2 x 50 mL) to remove the chlorobenzene. The methanolic phase was treated with 2 N hydrochloric acid (4 mL), producing a crystalline slurry. After one hour, the crystalline product was filtered, washed with methanol (5 mL), and dried at 60° in vacuo to give 2.23 g of the title compound. The X-ray powder diffraction pattern was identical to that reported in Table 3.

Table 3. X-ray Diffraction Pattern for Non-solvated Crystal Form.

	d-line spacing (Angstroms)	I/I ₀ (x100)
25	13.3864	71.31
	9.3598	33.16
	8.4625	2.08
	7.3888	7.57
	6.9907	5.80
30	6.6346	51.04
	6.1717	29.57
	5.9975	5.67
	5.9135	9.87
	5.6467	38.47
35	5.4773	10.54
	5.2994	4.74
	4.8680	4.03

-26-

	4.7910	5.98
	4.6614	57.50
	4.5052	5.75
	4.3701	9.03
5	4.2516	69.99
	4.2059	57.64
	4.1740	65.07
	4.0819	12.44
	3.9673	22.53
10	3.9318	100.00
	3.8775	9.07
	3.7096	33.38
	3.6561	21.65
	3.5576	3.36
15	3.5037	7.97
	3.4522	18.02
	3.4138	4.65
	3.2738	10.23
	3.1857	8.90
20	3.1333	6.24
	3.0831	9.43
	3.0025	12.13
	2.9437	4.96
	2.8642	7.70
25	2.7904	11.95
	2.7246	3.05
	2.6652	3.32
	2.5882	7.30

30 The amount of 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene hydrochloride present in the crystalline material is at least 95%.

35 This non-solvated crystalline form is particularly preferred for use in the manufacture of pharmaceutical compositions.

-27-

Example 1
Preparation of Amorphous Form

The amorphous material of the instant invention was prepared by dissolving 5 g of crystalline raloxifene hydrochloride in 300 ml of methanol and 22.5 ml of water. The amorphous material was recovered by spray drying the solution using a Buchi Model 190 Mini spray dryer under the following conditions: equilibrium inlet temperature: 84 °C; equilibrium outlet temperature: 60 °C; approx. spray rate: 2.5 ml/min.; aspirator setting: 20; air flow indicator: 500-600; atomization pressure: 35 psi. The process was completed in 2 hours and 10 minutes.

The material recovered after spray drying was observed under a polarized light microscope to determine if the product was birefringent. No birefringence was observed, and the material was considered to be amorphous.

Table 4 provides comparative solubility data in water between the crystalline and amorphous material:

20

Table 4

<u>Form</u>	<u>Solubility (mg/ml)</u>	<u>Temp., °C</u>
crystalline	0.08	room temperature
amorphous	42.2	room temperature
25 amorphous	120	37 °C *

* in sonicator

The amorphous material was also made as a complex with povidone, with ratios of raloxifene HCl to PVP as follows: 9:1, 3:1, and 1:1 (w/w). Complexes with hydroxypropyl- β -cyclodextrin (HPBCD) were also prepared, with a ratio of raloxifene HCl to HPBCD of from 1:1 to 1:4 (w/w). These were added to increase solubility and to prevent any potential crystallization of the material, for example, which may or may not occur when a supersaturated solution was prepared from the amorphous form. Some lots were recovered as crystalline

-28-

material, or the material was converted to a crystalline form within about a week or 10 days after preparation. The lots recovered as amorphous material which did not crystallize within about 10 days were observed to remain amorphous.

5

Example 2

Bioavailability Study

The extent and rate of absorption of the crystalline and amorphous forms of raloxifene HCl was determined. The two forms of the compound were formulated in a PEG vehicle as follows:

	<u>Component</u>	<u>% by weight</u>
	Polyethylene glycol 1450	70
	spray dried lactose	1.5
15	colloidal silicon dioxide	1.5
	polysorbate 80	2.0
	raloxifene HCl	25

Six dogs were dosed as follows: three dogs received the amorphous form, while three dogs received the crystalline form, both at a dose of 4mg/kg. Plasma levels of the two compound forms were then determined over a 10 hour period. The results from this study are provided in Figure 1. This study indicated greater bioavailability with the amorphous form.

The following discussions illustrate methods of use for the compounds of formula I in experimental models or in clinical studies. These examples are for the purposes of illustration and are not meant to be limiting in any way.

A. Osteoporosis:

Experimental models of postmenopausal osteoporosis are known in the art. Germane to this invention is the ovariectomized rat model which is provided in U.S. 5,393,763. The compounds of formula I would be active in this model and would demonstrate an effective treatment or prevention of bone loss due to the deprivation of estrogen.

An additional demonstration of the method of treating or preventing osteoporosis due to estrogen deprivation would

-29-

be as follows: One hundred patients would be chosen, who are healthy postmenopausal women, aged 45-60 and who would normally be considered candidates for estrogen replacement therapy. This includes women with an intact uterus, who
5 have had a last menstrual period more than six months, but less than six years. Patients excluded for the study would be those who have taken estrogens, progestins, or corticosteroids six months prior to the study or who have ever taken bis-phosphonates.

10 Fifty women (test group) would receive 15-80 mg of a compound of formula I, for example, Formulation 1 (above), per day. The other fifty women (control group) would receive a matched placebo per day. Both groups would receive calcium carbonate tablets (648 mg) per day. The
15 study is a double-blind design. Neither the investigators nor the patients would know to which group each patient is assigned.

A baseline examination of each patient includes quantitative measurement of urinary calcium, creatinine,
20 hydroxyproline, and pyridinoline crosslinks. Blood samples are measured for serum levels of osteocalcin and bone-specific alkaline phosphatase. Baseline measurements would also include a uterine examination and bone mineral density determination by photon absorptiometry.

25 The study would continue for six months, and each of the patients would be examined for changes in the above parameters. During the course of treatment, the patients in the treatment group would show a decreased change in the biochemical markers of bone resorption as compared to the
30 control group. Also, the treatment group would show little or no decrease in bone mineral density compared to the control group. Both groups would have similar uterine histology, indicating the compounds of formula I have little or no uterotrophic effects.

35 B. Hyperlipidemia:

Experimental models of postmenopausal hyperlipidemia are known in the art. Germane to this invention is the

-30-

ovariectomized rat model which is detailed in U.S.

5,464,845. Estrogenicity may further be assessed by evaluating the response of eosinophil infiltration into the uterus. A demonstration of the method of treating

5 hyperlipidemia due to estrogen deprivation would be as follows: One hundred patients would be chosen, who are healthy postmenopausal women, aged 45-60, and who would normally be considered candidates for estrogen replacement therapy. This would include women with an intact uterus,
10 who have not had a menstrual period for more than six months, but less than six years. Patients excluded for the study would be those who have taken estrogens, progestins, or corticosteroids.

Fifty women (test group) would receive 15-80 mg of a
15 compound of formula I, for example, using Formulation 1, per day. The other fifty women (control group) would receive a matched placebo per day. The study would be a double-blind design. Neither the investigators nor the patients would know to which group each patient is assigned.


20 A baseline examination of each patient would include serum determination of cholesterol and tri-glyceride levels. At the end of the study period (six months), each patient would have their serum lipid profile taken. Analysis of the data would confirm a lowering of the serum lipids, for
25 example, cholesterol and/or tri-glycerides, in the test group versus the control.

From the foregoing, it will be seen that this invention is one well adapted to attain all the ends hereinabove set forth together with advantages that are inherent to the
30 invention. It will be understood that certain features and subcombinations are of utility and may be employed without reference to other features and subcombinations. This is contemplated by and within the scope of the claims. Because many possible embodiments can be made of the invention
35 without departing from the scope thereof, it is to be understood that all matter herein set forth is to be interpreted as illustrative and not in a limiting sense.

-31-

We claim:

1. A compound of formula I



(I)

wherein:

R¹ and R³ are independently hydrogen, -CH₃, -CO(C₁-C₆ alkyl), or -COAr, wherein Ar is optionally substituted phenyl;


R² is selected from the group consisting of pyrrolidinyl, hexamethyleneimino, and piperidinyl; or a pharmaceutically acceptable salt or solvate thereof, in an amorphous form.

2. A compound according to Claim 1 wherein R¹ and R³ each are hydrogen, R² is piperidinyl, and the hydrochloride salt thereof.

3. A compound according to Claim 1 which is 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinylethoxy)benzoyl]benzo[b]thiophene.

-32-

4. A process for preparing an amorphous form of a compound of formula I



(I)

wherein:

R^1 and R^3 are independently hydrogen, $-CH_3$, $-CO(C_1-C_6 \text{ alkyl})$, or $-COAr$, wherein Ar is optionally substituted phenyl;

R^2 is selected from the group consisting of pyrrolidinyl, hexamethyleneimino, and piperidinyl; or a pharmaceutically acceptable salt or solvate thereof;
which comprises preparing a solution of a crystalline form of a compound of formula I in a suitable solvent, and then spray drying said solution to recover an amorphous form of a compound of formula I.

5. The process of claim 4 wherein said suitable solvent is a mixture of methanol and water.

6. The process of claim 4 wherein said spray drying is effected in the presence of an inert gas.

-33-

7. A pharmaceutical formulation comprising a compound according to Claim 1 in combination with a pharmaceutically acceptable carrier, diluent, or excipient.

8. A method for the inhibition of bone loss or bone resorption which comprises administering to a patient in need thereof a compound according to Claim 1.

9. The method according to Claim 8 wherein said bone loss or bone resorption occurs as a result of menopause or ovariectomy.

10. A method of inhibiting serum cholesterol levels which comprises administering to a patient in need thereof a compound according to Claim 1.

11. A method of alleviating the symptoms of post-menopausal syndrome which comprises administering to a patient in need thereof a compound according to Claim 1.

12. A method according to Claim 11 wherein said symptom of the post-menopausal syndrome pathological condition is osteoporosis.

13. A method according to Claim 11 wherein said symptom of the post-menopausal syndrome pathological condition is a cardiovascular disease.

14. A method according to Claim 13 wherein said cardiovascular disease is hyperlipidemia.

15. A method according to Claim 11 wherein said symptom of post-menopausal syndrome pathological condition is estrogen-dependent cancer.

16. A method according to Claim 15 wherein said estrogen-dependent cancer is breast or uterine cancer.

17. A method for alleviating the symptoms of uterine fibroid disease which comprises administering to a patient in need thereof a compound according to Claim 1.

18. A method for alleviating the symptoms of endometriosis which comprises administering to a patient in need thereof a compound according to Claim 1.

19. A method for alleviating the symptoms of aortal smooth muscle cell proliferation which comprises administering to a patient in need thereof a compound according to Claim 1.

20. A method for alleviating the symptoms of restenosis which comprises administering to a patient in need thereof a compound according to Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/14768**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 31/445; C07D 401/12

US CL :514/324; 546/281.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/324; 546/281.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ELKS, J. Dictionary of Drugs. London: Chapman and Hall. 1990, pages 1063 and 1064.	1-20, in part

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

23 OCTOBER 1997

Date of mailing of the international search report

17 NOV 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

R. W. RAMSUEY

Telephone No. (703) 306-4235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/14768

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-20, in part,
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Please See Extra Sheet.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JS92/14768

BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

2. Where no meaningful search could be carried out, specifically:

The multitude of variables and their permutations and combinations (e.g. R1,R2,Ar, the various methods of use,etc.) result in claimed subject matter that is so broad in scope that it is rendered virtually incomprehensible and thus no meaningful search can be given. Note also that the claimed subject matter lacks a significant structural element qualifying as the special technical feature that clearly defines a contribution over the art. The subject matter claimed contains an ethoxybenzoyl phenyl benzothiophene group which does not define a contribution over the prior art. Therefore, the first discernible invention as found in Example 1,(the compound therein, the pharmaceutical composition therewith, the method of preparation thereof and the method of inhibiting bone loss therewith) has been searched.